



# Cambridge International AS & A Level

CANDIDATE  
NAME


|  |
|--|
|  |
|--|

CENTRE  
NUMBER

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

CANDIDATE  
NUMBER

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|



**FURTHER MATHEMATICS**

**9231/32**

Paper 3 Further Mechanics

**October/November 2020**

**1 hour 30 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

## INSTRUCTIONS


- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ ms}^{-2}$ .

## INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.

1



A fixed smooth solid sphere has centre  $O$  and radius  $a$ . A particle of mass  $m$  is projected downwards with speed  $\sqrt{\frac{1}{6}ag}$  from the point  $A$  on the surface of the sphere, where  $OA$  makes an angle  $\alpha$  with the upward vertical through  $O$  (see diagram). The particle moves in part of a vertical circle on the surface of the sphere. It loses contact with the sphere at the point  $B$ , where  $OB$  makes an angle  $\beta$  with the upward vertical through  $O$ .

Given that  $\cos \alpha = \frac{2}{3}$ , find the value of  $\cos \beta$ . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



Two uniform smooth spheres  $A$  and  $B$  of equal radii have masses  $2m$  and  $m$  respectively. Sphere  $B$  is at rest on a smooth horizontal surface. Sphere  $A$  is moving on the surface with speed  $u$  and collides with  $B$ . Immediately before the collision, the direction of motion of  $A$  makes an angle  $\alpha$  with the line of centres of the spheres, where  $\tan \alpha = \frac{4}{3}$  (see diagram). The coefficient of restitution between the spheres is  $\frac{1}{3}$ .

Find the speed of  $A$  after the collision.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 An object consists of a uniform solid circular cone, of vertical height  $4r$  and radius  $3r$ , and a uniform solid cylinder, of height  $4r$  and radius  $3r$ . The circular base of the cone and one of the circular faces of the cylinder are joined together so that they coincide. The cone and the cylinder are made of the same material.

- (a) Find the distance of the centre of mass of the object from the end of the cylinder that is not attached to the cone. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(b)** Show that the object can rest in equilibrium with the curved surface of the cone in contact with a horizontal surface. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 A particle  $P$  of mass  $m$  is moving in a horizontal circle with angular speed  $\omega$  on the smooth inner surface of a hemispherical shell of radius  $r$ . The angle between the vertical and the normal reaction of the surface on  $P$  is  $\theta$ .

(a) Show that  $\cos \theta = \frac{g}{\omega^2 r}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The plane of the circular motion is at a height  $x$  above the lowest point of the shell. When the angular speed is doubled, the plane of the motion is at a height  $4x$  above the lowest point of the shell.

(b) Find  $x$  in terms of  $r$ .

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 A particle  $P$  is projected with speed  $u \text{ m s}^{-1}$  at an angle of  $\theta$  above the horizontal from a point  $O$  on a horizontal plane and moves freely under gravity. The horizontal and vertical displacements of  $P$  from  $O$  at a subsequent time  $t$  s are denoted by  $x$  m and  $y$  m respectively.

(a) Starting from the equation of the trajectory given in the List of formulae (MF19), show that

$$y = x \tan \theta - \frac{gx^2}{2u^2}(1 + \tan^2 \theta). \quad [1]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

When  $\theta = \tan^{-1} 2$ ,  $P$  passes through the point with coordinates (10, 16).

(b) Show that there is no value of  $\theta$  for which  $P$  can pass through the point with coordinates (18, 30). [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





6 One end of a light elastic string, of natural length  $a$  and modulus of elasticity  $k$ , is attached to a particle  $P$  of mass  $m$ . The other end of the string is attached to a fixed point  $Q$ . The particle  $P$  is projected vertically upwards from  $Q$ . When  $P$  is moving upwards and at a distance  $\frac{4}{3}a$  directly above  $Q$ , it has a speed  $\sqrt{2ga}$ . At this point, its acceleration is  $\frac{7}{3}g$  downwards.

Show that  $k = 4mg$  and find in terms of  $a$  the greatest height above  $Q$  reached by  $P$ . [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of horizontal dotted lines for writing.

7 A particle  $P$  of mass  $m$  kg moves in a horizontal straight line against a resistive force of magnitude  $mkv^2$  N, where  $v$  ms<sup>-1</sup> is the speed of  $P$  after it has moved a distance  $x$  m and  $k$  is a positive constant. The initial speed of  $P$  is  $u$  ms<sup>-1</sup>.

(a) Show that  $x = \frac{1}{k} \ln 2$  when  $v = \frac{1}{2}u$ . [4]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

Beginning at the instant when the speed of  $P$  is  $\frac{1}{2}u$ , an additional force acts on  $P$ . This force has magnitude  $\frac{5m}{v}$  N and acts in the direction of increasing  $x$ .

- (b) Show that when the speed of  $P$  has increased again to  $u \text{ m s}^{-1}$ , the total distance travelled by  $P$  is given by an expression of the form

$$\frac{1}{3k} \ln \left( \frac{A - ku^3}{B - ku^3} \right),$$

stating the values of the constants  $A$  and  $B$ .

[7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.