



## Cambridge International AS & A Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**MATHEMATICS**

**9709/41**

Paper 4 Mechanics

**October/November 2020**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ m s}^{-2}$ .

### INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **12** pages. Blank pages are indicated.

**1** A particle *B* of mass 5 kg is at rest on a smooth horizontal table. A particle *A* of mass 2.5 kg moves on the table with a speed of  $6 \text{ m s}^{-1}$  and collides directly with *B*. In the collision the two particles coalesce.

**(a)** Find the speed of the combined particle after the collision. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(b)** Find the loss of kinetic energy of the system due to the collision. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 A car of mass 1400 kg is moving along a straight horizontal road against a resistance of magnitude 350 N.

(a) Find, in kW, the rate at which the engine of the car is working when it is travelling at a constant speed of  $20 \text{ m s}^{-1}$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the acceleration of the car when its speed is  $20 \text{ m s}^{-1}$  and the engine is working at 15 kW. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....







Two particles of masses 0.8 kg and 0.2 kg are connected by a light inextensible string that passes over a fixed smooth pulley. The system is released from rest with both particles 0.5 m above a horizontal floor (see diagram). In the subsequent motion the 0.2 kg particle does not reach the pulley.

- (a) Show that the magnitude of the acceleration of the particles is  $6 \text{ m s}^{-2}$  and find the tension in the string. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....













