

Thursday 6 June 2019 – Afternoon

A Level Further Mathematics A

Y541/01 Pure Core 2

Time allowed: 1 hour 30 minutes



You must have:

- Printed Answer Booklet
- Formulae A Level Further Mathematics A

You may use:

• a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Write your answer to each question in the space provided in the Printed Answer **Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $gm s^{-2}$. Unless otherwise instructed, when a numerical value is needed, use g = 9.8.

INFORMATION

- The total mark for this paper is **75**.
- The marks for each question are shown in brackets [].
- You are reminded of the need for clear presentation in your answers.
- The Printed Answer Booklet consists of 16 pages. The Question Paper consists of 8 pages.

Answer all the questions.

1 In this question you must show detailed reasoning.

(a) By using partial fractions show that $\sum_{r=1}^{n} \frac{1}{r^2 + 3r + 2} = \frac{1}{2} - \frac{1}{n+2}$. [5]

(**b**) Hence determine the value of
$$\sum_{r=1}^{\infty} \frac{1}{r^2 + 3r + 2}$$
. [2]

2 (a) A plane
$$\Pi$$
 has the equation $\mathbf{r} \cdot \begin{pmatrix} 3 \\ 6 \\ -2 \end{pmatrix} = 15 \cdot C$ is the point (4, -5, 1).

Find the shortest distance between Π and *C*.

(b) Lines l_1 and l_2 have the following equations.

$$l_1: \mathbf{r} = \begin{pmatrix} 4\\3\\1 \end{pmatrix} + \lambda \begin{pmatrix} -2\\4\\-2 \end{pmatrix}$$
$$l_2: \mathbf{r} = \begin{pmatrix} 5\\2\\4 \end{pmatrix} + \mu \begin{pmatrix} 1\\-2\\1 \end{pmatrix}$$

Find, in exact form, the distance between l_1 and l_2 . [5]

[3]

3 In this question you must show detailed reasoning.

Show that
$$\int_{5}^{\infty} (x-1)^{-\frac{3}{2}} dx = 1.$$
 [5]

- 4 A 2-D transformation T is a shear which leaves the *y*-axis invariant and which transforms the object point (2, 1) to the image point (2, 9). A is the matrix which represents the transformation T.
 - (a) Find A. [3]
 - (b) By considering the determinant of A, explain why the area of a shape is invariant under T. [2]
- 5 A particle of mass 2kg moves along the x-axis. At time t seconds the velocity of the particle is $v \text{ ms}^{-1}$.

The particle is subject to two forces.

- One acts in the positive x-direction with magnitude $\frac{1}{2}t$ N.
- One acts in the negative *x*-direction with magnitude *v* N.
- (a) Show that the motion of the particle can be modelled by the differential equation

$$\frac{\mathrm{d}v}{\mathrm{d}t} + \frac{1}{2}v = \frac{1}{4}t.$$
[1]

The particle is at rest when t = 0.

- (b) Find v in terms of t. [5]
- (c) Find the velocity of the particle when t = 2. [1]

When t = 2 the force acting in the **positive** *x*-direction is replaced by a constant force of magnitude $\frac{1}{2}$ N in the same direction.

- (d) Refine the differential equation given in part (a) to model the motion for $t \ge 2$. [1]
- (e) Use the refined model from part (d) to find an exact expression for v in terms of t for $t \ge 2$.

[3]

6 *A* is a fixed point on a smooth horizontal surface. A particle *P* is initially held at *A* and released from rest.

It subsequently performs simple harmonic motion in a straight line on the surface. After its release it is next at rest after 0.2 seconds at point B whose displacement is 0.2 m from A. The point M is halfway between A and B.

The displacement of P from M at time t seconds after release is denoted by x m.

- (a) On the axes provided in the Printed Answer Booklet, sketch a graph of x against t for $0 \le t \le 0.4$. [4]
- (b) Find the displacement of *P* from *M* at 0.75 seconds after release. [2]
- 7 In an Argand diagram the points representing the numbers 2 + 3i and 1 i are two adjacent vertices of a square, *S*.
 - (a) Find the area of S. [3]
 - (b) Find all the possible pairs of numbers represented by the other two vertices of *S*. [4]

8 In this question you must show detailed reasoning.


(a) By writing $\sin\theta$ in terms of $e^{i\theta}$ and $e^{-i\theta}$ show that

$$\sin^{6}\theta = \frac{1}{32}(10 - 15\cos 2\theta + 6\cos 4\theta - \cos 6\theta).$$
 [5]

(**b**) Hence show that
$$\sin \frac{1}{8}\pi = \frac{1}{2} \sqrt[6]{20 - 14\sqrt{2}}$$
. [3]

9 In this question you must show detailed reasoning.

The diagram below shows the curve $r = \sqrt{\sin\theta} e^{\frac{1}{3}\cos\theta}$ for $0 \le \theta \le \pi$.



- (a) Find the exact area enclosed by the curve. [4] (b) Show that the greatest value of *r* on the curve is $\sqrt{\frac{\sqrt{3}}{2}}e^{\frac{1}{6}}$. [7]
- 10 (a) Use differentiation to find the first two non-zero terms of the Maclaurin expansion of $\ln(\frac{1}{2} + \cos x)$. [4]

(b) By considering the root of the equation $\ln\left(\frac{1}{2} + \cos x\right) = 0$ deduce that $\pi \approx 3\sqrt{3\ln\left(\frac{3}{2}\right)}$. [3]

END OF QUESTION PAPER

BLANK PAGE

BLANK PAGE



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.