



Pearson
Edexcel

Mark Scheme (Results)

Autumn 2020

Pearson Edexcel GCE Further Mathematics
AS Further Mechanics 2 Paper 8FM0_26

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Autumn 2020

Publications Code 8FM0_26_2010_MS

All the material in this publication is copyright

© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the last candidate in exactly the same way as they mark the first.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification/indicative content will not be exhaustive.

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 40.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 5. Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response.
If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is the most complete.
 6. Ignore wrong working or incorrect statements following a correct answer.
 7. Mark schemes will firstly show the solution judged to be the most common response expected from candidates. Where appropriate, alternatives answers are provided in the notes. If examiners are not sure if an answer is acceptable, they will check the mark scheme to see if an alternative answer is given for the method used.

General Principles for Mechanics Marking

(But note that specific mark schemes may sometimes override these general principles)

- Rules for M marks: correct no. of terms; dimensionally correct; all terms that need resolving (i.e. multiplied by cos or sin) are resolved.
- Omission or extra g in a resolution is an accuracy error not method error.
- Omission of mass from a resolution is a method error.
- Omission of a length from a moments equation is a method error.
- Omission of units or incorrect units is not (usually) counted as an accuracy error.
- dM indicates a dependent method mark i.e. one that can only be awarded if a previous specified method mark has been awarded.
- Any numerical answer which comes from use of $g = 9.8$ should be given to 2 or 3 SF.
- Use of $g = 9.81$ should be penalised once per (complete) question.

N.B. Over-accuracy or under-accuracy of correct answers should only be penalised *once* per complete question. However, premature approximation should be penalised every time it occurs.

- Marks must be entered in the same order as they appear on the mark scheme.
- In all cases, if the candidate clearly labels their working under a particular part of a question i.e. (a) or (b) or (c),.....then that working can only score marks for that part of the question.
- Accept column vectors in all cases.
- Misreads – if a misread does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, bearing in mind that after a misread, the subsequent A marks affected are treated as A ft
- Mechanics Abbreviations
 - M(A) Taking moments about A
 - N2L Newton's Second Law (Equation of Motion)
 - NEL Newton's Experimental Law (Newton's Law of Impact)
 - HL Hooke's Law
 - SHM Simple harmonic motion
 - PCLM Principle of conservation of linear momentum
 - RHS, LHS Right hand side, left hand side

Question	Scheme	Marks	AOs
1(a)	$\frac{1}{2}a$	B1	1.1b
	Loaded lamina has a mass distribution which is symmetrical about the perpendicular bisector of AD	B1	2.4
		(2)	
(b)	Moments about AD	M1	3.1a
	$6ma + m.2a = 12m\bar{x}$	A1	1.1b
	$\bar{x} = \frac{2a}{3}$ *	A1*	2.2a
		(3)	
(c)	Moments about AB	M1	3.1a
	$kma + 12m.\frac{1}{2}a = (k+12)m\bar{y}$	A1	1.1b
		A1	1.1b
	$\bar{y} = \frac{(k+6)a}{(k+12)}$ *	A1*	2.2a
		(4)	
(d)	Moments about AD	M1	3.1a
	$\bar{x}_1 = \frac{8a}{(k+12)}$	A1	1.1b
	Use of $\tan \alpha = \frac{\bar{y}}{\bar{x}_1}$	M1	1.1b
	$\frac{3}{2} = \frac{\frac{(k+6)a}{(k+12)}}{\frac{8a}{(k+12)}}$	A1	1.1b
	Solve for k	M1	1.1b
	$k = 6$	A1	1.1b
	SC: For use of $(\tan \alpha =) \frac{\text{their } \bar{y}}{\text{their } \bar{x}} = \frac{3}{2}$, M1A1M0A0M0A0		
		(6)	
			(15 marks)

Notes

(a)

B1: cao

B1: clear explanation

(b)

M1: Correct no. of terms and dimensionally correct (allow cancelled $m's$)

A1: A correct equation

A1*: Correctly obtained printed answer

(c)

M1: Correct no. of terms and dimensionally correct (allow cancelled $m's$)

A1: Correct equation with one error

A1: Correct equation

A1*: Correctly obtained printed answer

(d)

M1: Correct no. of terms and dimensionally correct (allow cancelled $m's$)

A1: Correct distance

M1: Correct use of tan but allow reciprocal

A1: Correct equation

M1: Solve for k

A1: cao

Question	Scheme	Marks	AOs
2(a)	$(a\sqrt{3})^2 + (3a - AP)^2 = AP^2$	M1	1.1b
	$AP = 2a$ *	A1*	1.1b
		(2)	
(b)	Equation of motion horizontally	M1	3.1b
	$T + T \times \frac{1}{2} = \frac{mU^2}{a}$	A1	1.1b
		A1	1.1b
	$T = \frac{2mU^2}{3a}$	A1	2.2a
	(4)		
(c)	Resolving vertically	M1	3.1b
	$R + T \times \frac{\sqrt{3}}{2} = mg$	A1	1.1b
	On the table $\Rightarrow R > 0$	M1	2.1
	$mg - \frac{2mU^2\sqrt{3}}{3a \times 2} > 0$	A1	1.1b
	$U^2 < ag\sqrt{3}$ *	A1*	2.2a
		(5)	
(d)	Bead would lift off the table	B1	2.4
		(1)	
(e)	Tension would vary along the string	B1	3.5b
		(1)	
			(13 marks)

Notes

(a)

M1: Use of Pythagoras $3a^2 + 9a^2 - 6a \times AP + AP^2 = AP^2 \Rightarrow 6a \times AP = 12a^2$

A1: $AP = 2a$. GIVEN ANSWER

(b)

M1: Use of horizontal equation to solve the problem, with correct no. of terms etc

A1: Equation with at most one error

A1: Correct equation

A1: Correct answer

(c)

M1: Use of vertical resolution to solve the problem, with correct no. of terms etc

A1: Correct equation

M1: Use of $R > 0$

A1: Correct inequality

A1*: Correctly obtained given answer

(d)

B1: Clear comment

(e)

B1: Clear explanation

Question	Scheme	Marks	AOs
3(a)	Use of initial condition to find p	M1	3.1b
	$t = 0, v = 0, \text{ acceleration} = 3 \Rightarrow p = 3$	A1	1.1b
	Use $v = 4, \text{ acceleration} = \frac{1}{2}$	M1	1.1b
	$q = -\frac{5}{8}$	A1	1.1b
	Use acceleration = $\frac{dv}{dt}$ and rearrange	M1	1.1b
	$8\frac{dv}{dt} = (24 - 5v) *$	A1*	2.2a
		(6)	
(b)	Separate the variables and integrate	M1	3.1b
	$8\int \frac{dv}{(24 - 5v)} = \int dt$	A1	1.1b
	$-\frac{8}{5}\ln(24 - 5v) = t + C$	A1	1.1b
	Use $t = 0, v = 0$ to give $C = -\frac{8}{5}\ln 24$	M1	1.1b
	Substitute $v = 4$ and find and simplify T	M1	1.1b
	$T = \frac{8}{5}\ln 6$	A1	1.1b
		(6)	
(12 marks)			

Notes
<p>(a) M1: Use initial conditions A1: cao M1: Use second condition A1: cao M1: Use appropriate derivative and rearrange A1*: Correct given answer</p>
<p>(b) M1: Separate the variables and integrate A1: Correct integration (C not required) M1: Use of limits or initial conditions to find C M1: Use $v = 4$ to find T A1: Correct answer (single log)</p>

