

Cambridge International AS & A Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

0123456789

MATHEMATICS 9709/01

Paper 1 Pure Mathematics 1

For examination from 2020

SPECIMEN PAPER

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has 22 pages. Blank pages are indicated.

© UCLES 2017 [Turn over

BLANK PAGE

1 The following points

A(0, 1), B(1, 6), C(1.5, 7.75), D(1.9, 8.79) and E(2, 9)

lie on the curve y = f(x). The table below shows the gradients of the chords AE and BE.

Chord	AE	BE	CE	DE
Gradient of chord	4	3		

(a)	Complete the table to show the gradients of <i>CE</i> and <i>DE</i> .	[2]
(b)	State what the values in the table indicate about the value of $f'(2)$.	[1]

 $f: x \mapsto 3x + 2, \quad x \in \mathbb{R},$

2	Functions	f	and	ø	are	defined	þν
_	1 uncuons		unu	~	ui C	actifica	$\boldsymbol{\nu}$

$g: x \mapsto 4x - 12,$	$x \in \mathbb{R}$.
Solve the equation $f^{-1}(x) = gf(x)$.	[4]

Find the value of n .	

١	
	Find $f(x)$.

Find and simplify the equation of the translated curve.	
	[2]
(b) The graph of $y = f(x)$ is transformed to the graph of $y = 3f(-x)$.	
Describe fully the two single transformations which have been contransformation.	mbined to give the resulting [3]

••••••
•••••
 •••••

 •••••
 •••••
•••••
••••••
••••••
•••••
••••••
 •••••
 •••••
 •••••


	$6\cos^2 x - \cos x - 1 = 0. ag{3}$
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	
•	

•••••
••••••
••••••
••••••
 •••••
 •••••
•••••
•••••
•••••

o	A 1	12
8	A curve has equation $y =$	$\overline{3-2x}$

(a)	Find $\frac{dy}{dx}$.	[2]
	oint moves along this curve. As the point passes through A , the x -co units per second and the y -coordinate is increasing at a rate of 0.4	
(b)	Find the possible x -coordinates of A .	[4

9



The diagram shows a circle with centre A and radius r. Diameters CAD and BAE are perpendicular to each other. A larger circle has centre B and passes through C and D.

(a)	Show that the radius of the larger circle is $r\sqrt{2}$.	[1]
(b)	Find the area of the shaded region in terms of r .	[6]
		•••••
		•••••
		•••••
		•••••
		•••••
		•••••

,	The	circle $x + y + 4x - 2y - 20 = 0$ has centre C and passes through points A and B.	
	(a)	State the coordinates of <i>C</i> .	[1]
	It is	given that the midpoint, D, of AB has coordinates $(1\frac{1}{2}, 1\frac{1}{2})$.	
		Find the equation of AB, giving your answer in the form $y = mx + c$.	[4]
	(D)	That the equation of MD , giving your answer in the form $y = mx + c$.	נדן
			•••••
			•••••
			•••••
			•••••
			•••••
			•••••
			•••••


•		
•		•••••
•		
•		•••••
•		
•		••••••••••
_	Find by coloulation the a goodinates of A and D	
ŀ	Find, by calculation, the x -coordinates of A and B .	
•		
•		••••••••••
•		
•		
		•••••
•		
•		
•••		
• • • • • • • • • • • • • • • • • • • •		
••		
• •		
•		

11 The function f is defined, for $x \in \mathbb{R}$, by $f: x \mapsto x^2 + ax + b$, where a and b are constants.

(a)	It is given that $a = 6$ and $b = -8$.	
	Find the range of f.	[3]
(b)	It is given instead that $a = 5$ and that the roots of the equation $f(x) = 0$ constant.	are k and $-2k$, where k is a
	Find the values of b and k .	[3]

	•••••
	•••••
Show that if the equation $f(x + a) = a$ has no real roots then $a^2 < 4(b - a)$.	[3]
	••••••
	••••••
	•••••
	now that if the equation $f(x + a) = a$ has no real roots then $a^2 < 4(b - a)$.

12



The diagram shows the curve with equation $y = x(x - 2)^2$. The minimum point on the curve has coordinates (a, 0) and the x-coordinate of the maximum point is b, where a and b are constants.

(a)	State the value of a.	[1]
(b)	Calculate the value of b .	[4]
		••••••

The gradient, $\frac{dy}{dx}$, of the curve has a minimum value m .	
	Γ
The gradient, $\frac{dy}{dx}$, of the curve has a minimum value m . Calculate the value of m .	[-
	[-
	[
	[.
	[-
	[

Additional page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s must be clearly shown.			

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.