Please check the examination de	tails below before ente	ring your candidate information
Candidate surname		Other names
Pearson Edexcel Level 3 GCE	Centre Number	Candidate Number
Thursday 16	<b>May 20</b>	19
Afternoon	Paper Re	eference <b>8FM0-26</b>
Further Mathe Advanced Subsidiary Further Mathematics options 26: Further Mechanics 2 (Part of option J only)		
You must have: Mathematical Formulae and St	atistical Tables (Gre	een), calculator

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of g is required, take  $g = 9.8 \text{ m s}^{-2}$  and give your answer to either 2 significant figures or 3 significant figures.

## **Information**

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 40. There are 4 questions.
- The marks for each question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶










Figure 1

Five identical uniform rods are joined together to form the rigid framework ABCD shown in Figure 1. Each rod has weight W and length 4a. The points A, B, C and D all lie in the same plane.

The centre of mass of the framework is at the point G.

(a) Explain why G is the midpoint of AC.

**(1)** 

The framework is suspended from the ceiling by two vertical light inextensible strings. One string is attached to the framework at A and the other string is attached to the framework at B. The framework hangs freely in equilibrium with AB horizontal.

- (b) Find
  - (i) the tension in the string attached at A,
  - (ii) the tension in the string attached at B.

**(4)** 

A particle of weight kW is now attached to the framework at D and a particle of weight 2kW is now attached to the framework at C. The framework remains in equilibrium with AB horizontal and the strings vertical.

Either string will break if the tension in it exceeds 6W.

(c) Find the greatest possible value of k.

**(4)** 


Question 1 continued



Question 1 continued	

Question 1 continued	
(Total for Question 1	is 9 marks)



**2.** A car moves in a straight line along a horizontal road. The car is modelled as a particle. At time t seconds, where  $t \ge 0$ , the speed of the car is  $v \text{ m s}^{-1}$ 

At the instant when t = 0, the car passes through the point A with speed 2 m s<sup>-1</sup>

The acceleration,  $a \text{ m s}^{-2}$ , of the car is modelled by

$$a = \frac{4}{2 + v}$$

in the direction of motion of the car.

(a) Use algebraic integration to show that  $v = \sqrt{8t + 16} - 2$ 

**(6)** 

At the instant when the car passes through the point B, the speed of the car is  $4 \,\mathrm{m\,s}^{-1}$ 

(b) Use algebraic integration to find the distance AB.

**(6)** 


Question 2 continued



Question 2 continued	

Question 2 continued	
/ID ( 3.0	O
(Total for	Question 2 is 12 marks)



**3.** A light inextensible string has length 8a. One end of the string is attached to a fixed point A and the other end of the string is attached to a fixed point B, with A vertically above B and AB = 4a. A small ball of mass m is attached to a point P on the string, where AP = 5a.

The ball moves in a horizontal circle with constant speed v, with both AP and BP taut.

The string will break if the tension in it exceeds  $\frac{3mg}{2}$ 

By modelling the ball as a particle and assuming the string does not break,

(a) show that 
$$\frac{9ag}{4} < v^2 \leqslant \frac{27ag}{4}$$

**(7**)

(b) find the least possible time needed for the ball to make one complete revolution.

**(2)** 

Question 3 continued



Question 3 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_

Question 3 continued	
(Te	otal for Question 3 is 9 marks)






Figure 2

The uniform triangular lamina ABCDE is such that angle  $CEA = 90^{\circ}$ , CE = 9a and EA = 6a. The point D lies on CE, with DE = 3a. The point B on CA is such that DB is parallel to EA and DB = 4a. The triangular lamina is folded along the line DB to form the folded lamina ABDECF, as shown in Figure 2.

The distance of the centre of mass of the triangular lamina from DC is  $d_1$ 

The distance of the centre of mass of the folded lamina from DC is  $d_{\gamma}$ 

(a) Explain why 
$$d_1 = d_2$$
 (1)

The folded lamina is freely suspended from B and hangs in equilibrium with BA inclined at an angle  $\alpha$  to the downward vertical through B.

(b) Find, to the nearest degree, the size of angle  $\alpha$ .



Question 4 continued



Question 4 continued	

Question 4 continued



Question 4 continued	

Question 4 continued



nestion 4 continued	
	(Total for Question 4 is 10 marks)
	TOTAL FOR FURTHER MECHANICS 2 IS 40 MARKS

