Please check the examination de	etails below before entering	your candidate information
Candidate surname	Ot	her names
Pearson Edexcel Level 3 GCE	Centre Number	Candidate Number
Thursday 14	May 202	0
Afternoon	Paper Refe	rence 8FM0/22
Further Mathe Advanced Subsidiary Further Mathematics o 22: Further Pure Mathe (Part of option A only)	ptions	
You must have: Mathematical Formulae and St	atistical Tables (Greer	Total Marks

Candidates may use any calculator allowed by Pearson regulations.
Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 40. There are 5 questions.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1.	The set $G = \{1, 3, 7, 9, 11, 13, 17, 19\}$ under the binary operation of multiplication modulo 20 forms a group.	
	(a) Find the inverse of each element of G.	(3)
	(b) Find the order of each element of G.	(3)
	(c) Find a subgroup of G of order 4	(1)
	(d) Explain how the subgroup you found in part (c) satisfies Lagrange's theorem.	(1)

Question 1 continued



Question 1 continued

Question 1 continued	
	Total for Question 1 is 2 mayles)
	Total for Question 1 is 8 marks)



2.	The highest common factor of 963 and 657 is c .	
	(a) Use the Euclidean algorithm to find the value of c .	(3)
	(b) Hence find integers a and b such that	
	963a + 657b = c	(3)

Question 2 continued



Question 2 continued

Question 2 continued
(Total for Question 2 is 6 marks)



3. (i)

$$\mathbf{A} = \begin{pmatrix} 1 & -2 \\ 1 & 4 \end{pmatrix}$$

(a) Show that the characteristic equation for **A** is $\lambda^2 - 5\lambda + 6 = 0$

(2)

(b) Use the Cayley-Hamilton theorem to find integers p and q such that

$$\mathbf{A}^3 = p\mathbf{A} + q\mathbf{I} \tag{3}$$

(ii) Given that the 2×2 matrix **M** has eigenvalues -1 + i and -1 - i,

with eigenvectors
$$\begin{pmatrix} 1 \\ 2-i \end{pmatrix}$$
 and $\begin{pmatrix} 1 \\ 2+i \end{pmatrix}$ respectively, find the matrix \mathbf{M} .

(5)

Question 3 continued	



Question 3 continued

Question 3 continued	
	Total for Question 3 is 10 marks)
	Total for Question 3 is to marks)



4. Sam borrows £10000 from a bank to pay for an extension to his house. The bank charges 5% annual interest on the portion of the loan yet to be repaid. Immediately after the interest has been added at the end of each year and before the start of the next year, Sam pays the bank a fixed amount, £F.

Given that $\pounds A_n$ (where $A_n \ge 0$) is the amount owed at the start of year n,

(a) write down an expression for A_{n+1} in terms of A_n and F,

(1)

(b) prove, by induction that, for $n \ge 1$

$$A_n = (10\,000 - 20F)1.05^{n-1} + 20F$$

(5)

(c) Find the smallest value of *F* for which Sam can repay all of the loan by the start of year 16.

(4)

Question 4 continued



Question 4 continued

Question 4 continued	
(TD 4.1	for Overtion A is 10 mag-l
(10tal :	for Question 4 is 10 marks)






Figure 1

Figure 1 shows an Argand diagram.

The set of points, A, that lies within the shaded region, including its boundaries, is defined by

$$A = \{z : p \leqslant \arg(z) \leqslant q\} \cap \{z : |z| \leqslant r\}$$

where p, q and r are positive constants.

(a) Write down the values of p, q and r.

(2)

Given that $w = -2\sqrt{3} + 2i$ and $z \in A$,

(b) find the maximum value of $|w-z|^2$ giving your answer in an exact simplified form.

(4)

Question 5 continued



Question 5 continued	
	(Total for Question 5 is 6 marks)
TOTAL FOR FURTHER PURE MATHEMATICS 2 IS 40 MARKS	

